



CLINICAL AND LABORATORY CHARACTERISTICS OF CHRONIC OSTEOMYELITIS IN CHILDREN

Berdiyaroova Sh. Sh.

Berdiyaroova Shokhida Shukurullaevna - Assistant; Samarkand State Medical University. Department of Clinical and Laboratory Diagnostics
(+99893) 358 13 41. berdiyarovashohida@gmail.com,

Yusupov Sh. A.

Yusupov Shukhrat Abdurasulovich - Associate Professor, Doctor of Medical Sciences, Samarkand State Medical University. Department of Pediatric Surgery
№ 1 (+99891)548 16 13. shuchrat_66@mail.ru

Yusupova N. A.

Yusupova Nargiza Abdikodirovna - Assistant, Samarkand State Medical University. Department of Clinical and Laboratory Diagnostics
(+99888) 406 44 48. nargiza-yusupova-87@mail.ru

Murtazaeva N. K.

Murtazaeva Nasiba Komilzhonovna - Assistant; Samarkand State Medical University. Department of Biological Chemistry (+99893) 353 81 55
murtazayevanasiba81@gmail.com.

Abstract

The publication summarizes various classical and modern aspects of acute and chronic hematogenous osteomyelitis in children. The results of our own observations and laboratory, instrumental, immunological studies carried out in a large contingent of patients with various forms of hematogenous osteomyelitis, as well as the results of various methods of surgical treatment; issues of etiology, pathogenesis, differential diagnosis and complex staged treatment of children were discussed; the original ideas about the modern classification of these actual diseases were expressed; there are six illustrative clinical examples from our own observations. The publication is intended for doctors of pediatric surgeons and specialists in related specialties.

Keywords: hemotogen osteomyelitis, bone tissue, collagen, forecasting, rehabilitation, children.





Introduction

The use of laboratory research methods in practice First of all, the study of blood is a laboratory method for detecting a bacterial pathogen. But this method is not so relevant for the chronic form of the disease that has developed in the human body. The inflammatory process causes leukocytosis, which in the acute phase, at the height of the disease, can reach 30 and above. Indicators such as ESR and C-reactive protein detect infection and show the severity of the disease. Often, during the examination, it is necessary to differentiate the disease with others, no less serious, such as post-traumatic periostitis, erysipelas, rheumatism, tuberculosis, neoplasms and actinomycosis of bone tissue. Methods of laboratory research in case of damage to the joints: The use of instrumental techniques for diagnosis. These methods include microbiological and radiological diagnostics. In order to correctly diagnose and determine the cause of the disease, it is necessary to investigate how pathogenic microorganisms are released in those places where the bone is damaged, as well as from the joint fluid and blood. Difficulties in the examination arise in patients with a developed hematogenous form, since a positive culture is detected in less than 50% of situations. One of the well-known studies is densitometry, which allows you to understand how bones are prone to fracture and destruction. Basic forms of microbiological research. This type of diagnosis has several ways to identify pathogens. Brief description of each of them: Bacterioscopy. Taking a smear from the mucous membrane in order to assess the state of the microflora. The preparation of the analysis takes place within a few days, and its decoding is carried out only by a doctor. Examination of the contents separated from the fistula during its formation. It may have purulent contents, as a result of which the diagnosis is difficult.

Serological Methods

Able to detect antigens and antibodies and include several reactions after a visual observation of the site of joint damage. Such methods allow not only to assess and detect the true causative agent of the infection, but also to consider the state of the microflora of the mucous membranes. Features of x-ray examination X-ray diagnostics is one of the most common methods for any type of osteomyelitis. It is based on the preparation of images of the affected limb, which are performed in two projections. In this way, it is possible to determine exactly where the infection is localized, how pronounced the symptoms are, as well as its prevalence in the tissues. In a child, manifestations of a developing disease can be noticed already on the fifth day, in adults - on the fifteenth. X-ray can be done several times during treatment to track the positive or negative dynamics. One of the disappointing indicators of the development





of the disease is the spread of the periosteal reaction to the previously healthy periosteum. X-ray osteomyelitic signs include: by the beginning of the second week, the line between compact and spongy substance disappears; bone fractures become oval or round; the relief of the periosteum thickens and changes; sequesters are noticeable by the end of the first month. An X-ray image allows you to visually study the state of the skeletal system, to prescribe medical procedures on-line, excluding the transition of the disease from an acute form to a chronic one. Especially often it is used in the study of diseases of the teeth. The importance of computed tomography CT is most often used in the chronic form, the occurrence of recurrent moments. It involves the image of a limb on a computer, allows you to evaluate not only the qualitative state of the bones, but also quantitative. This method makes it possible to detect sequesters of soft tissue areas that are affected and for a long period of time are not replaced by new growths, and purulent stagnation, if left untreated, leading to sepsis. Tomography makes it possible to build tactics for a phased surgical intervention if there are indications for this process. The effectiveness of fistulography in determining therapy. This technique can be attributed to the X-ray diagnostic method. Fistulography is a technique that involves the introduction of a certain substance into the fistula, and then fluoroscopy is performed in this area. Before preparing everything necessary for the formation of a fistulogram, an x-ray is preliminarily performed to analyze the cavity under study before the procedure begins. At the next stage, the edges of the fistula opening are lubricated with a solution of iodine with alcohol and the contents are taken, and then a contrast agent, prepared and warmed up in advance, is injected. When the fistula passages are completely filled, the cannula is taken out and the hole is sealed with a plaster. Then, within the framework of radiology, it is important to again conduct an examination using images, where it will be revealed how the fistulous passages are located. As a result, after the entire procedure, the fistula is released from the previously introduced content, the affected parts of the joints are processed and washed. The productivity of magnetic resonance imaging in determining the disease MRI allows you to determine in percentage terms how much the bone parts are affected by the infection. When carrying out this method, the doctor sees where there are still healthy tissues, and where damaged areas that look swollen are localized. According to experts, it is more effective than scintigraphy - the introduction of radioactive isotopes and the analysis of a two-dimensional image during radiation. Positive practice has also been recorded in the detection of sequesters. MRI is available at any stage of the course: when there are only prerequisites for pathology, when it is already developing, and even in severe cases, when the disease can be complicated. The use of scintigraphy as a method of rapid detection of infection It allows you to





explore the metabolism of the tissue system. It is carried out using radiation equipment (radionuclide diagnostics). A specialized substance is injected intravenously into the bone department, which tends to linger and accumulate in the bones. After that, the radiation is fixed by the detectors of the device that captures the signal. The introduction of the drug is accompanied by the use of 1 liter of water: thus, the accumulation of the contents is better and the radiation load on the body is reduced. Constructiveness in the use of ultrasound. This technique involves the use of equipment that performs ultrasound. Previously, among doctors it was believed that it is effective only in case of damage to soft tissues and cavities that contain fluid. Recently, it has been used in the examination of the hip joints, spine, and the detection of fractures of tubular bones. Ultrasound is used as a diagnosis of osteomyelitis of the upper or lower jaw, especially in children who are acutely experiencing inflammatory processes. Conclusion The effectiveness of each of the methods is confirmed by many years of practice of use by doctors actively involved in the diagnosis of osteomyelitis. They are aimed at early detection of the disease using various tests, tests, maintaining clinical protocols, studying tables and visual features of the manifestation of symptoms, reading analyzes at the biological level. If the development of the disease has already occurred, these techniques are aimed at shaping the doctor's understanding of the patient's health status, the clinical diagnostic picture of the signs, and his prospects in the process of the upcoming treatment. When a person has already received recommendations and frequent painful sensations in the joints cease to bother, these methods can help to identify the causes that entail not only primary symptoms, but also chronic relapses.

References

1. Akjigitov G.N.A., Gallayev M. A., Saxautov V.G., Yudin Ya.B. Osteomiyelit . M:Meditsina, 1986. 207 s.
2. Fedotov V.K. The system of diagnostic and treatment-rehabilitation measures for chronic osteomyelitis in children: Ph.D. dis. Dr. med. Sciences. M., 1992. 32 p.
3. Ali-Zade Ch.A. Treatment and prevention of recurrence of chronic hematogenous osteomyelitis in pediatric patients // Materials of Symp. det. traumatologists-orthopedists of Russia. SPb., 2003. S. 33-34.
4. Vengerovsky I. S. Osteomyelitis in children. Moscow: Medicine, 1964. 270 p.
5. Roshal L.M. Chronic osteomyelitis // Purulent surgery of children: a guide for physicians. L.: Medicine, 1991. S. 229-231.





6. Nabieva F.S., Dushanova G.A., Bobokulov O.O. Importance of enzyme immunoassay in the diagnosis of infectious diseases Vestnik nauki i obrazovaniya. No. 4(107). Part 1, 2021. 54-56 p.
7. Abdikadirova Nargiza Yusupovna, Berdiyaroova Shohida Shukuralievna, Yulaeva Irina Andreevna, Shodmanova Zebiniso Rahimovna. Clinical and Laboratory Parameters in Children with Urolithiasis and the Quality of Laboratory Tests at the Stage of Stationary. Treatment Annals of R.S.C.B., ISSN: 1583-6258. Vol. 25. Issue 1, 2021.
8. Hematological characteristics of risk factors and evaluation of prognosis in COVID-19. Yusupova N.A., Berdiyaroova Sh.Sh., Yulaeva I.A. Vesnik nauka I obrazovaniya., 2021. No. 5 (108). Part
9. Nargiza Yusupova, Oripov Firdavs. Energy drinks. The composition of energy drinks and the effect on the body of their individual components// Tematics journal of Microbiology. Vol-6-Issue-1-2022. P.24-35.
10. Berdiyaroova Sh.Sh., Yusupova N.A., Murtazaeva N.K., Ibragimova N.S. Clinical and laboratory features of chronic hematogenic osteomyelitis// TJM - Tematics journal of Microbiology ISSN 2277-2952 Vol-6-Issue-1-2022, P. 36 – 43.
11. Kudratova Z.E., Mukhammadieva L.A., Kuvandikov G.B. // Features of the etiopathogenesis of obstructive bronchitis and laryngotracheitis caused by atypical microflora / Achievements of science and education, No. 6 (77) 2020. P. 71-72.
12. Ibragimova N.S., Ibragimov B.F., Yusupova N.A., Yulaeva I.A. Course, complications and outcomes of postponed pregnancy// European science, 50-54, 2021.

