



REACTIVITY OF THE HYPOTHALAMIC-PITUITARY GONADAL SYSTEM AFTER 10 MINUTES OF CLINICAL DEATH

Karabaev Aminjon Gadaevich

M D, Associate Professor, Head of the Department of Physiology of
Samarkand State Medical University, Samarkand.Uzbekistan.

Mail: karabaev.aminion@bk.ru ORCID: 0000-0002-3355-0741.

Kim Diana Vladislavna

Assistant of the Department of Physiology of Samarkand
State Medical University, Samarkand.Uzbekistan.

Annotation

When modeling a 10-minute clinical death, changes in the reproductive system were studied in white mature male rats with a weight of 180-220 g. Clinical death was modeled according to the method of V. G. Korpachev (1982). The reproductive state in intact and experimental rats was studied using morphological, morphometric, histochemical, cytophotometric biochemical research methods. At the same time, the reproductive system in index rats is in a state of moderate functioning to highlight the synthesis and secretion of testosterone.

During the period of clinical death, there is an increase in the functional activity of d-basophil cells of the adenohypophysis, the content of the Lethinating hormone provided an increase in testosterone secretion

Keywords: Clinical death, endocrine system, adenohypophysis, follicle-stimulating hormone, lutinizing hormone, testosterone.

Relevance

A number of scientific studies are being conducted all over the world to improve the assessment of endocrine system disorders between anabolic, catabolic and kniteche systems under the influence of various extreme factors. [Zarechnova N.N., Slynko T.N., 2018; Yakimov I.A., Loginova E.S., 2017; Gadaevich K. A. et al. 2021; Karabayev A. G., 2020; Karabayev A. G., Isroilov R. I., 2020].

We know that when exposed to extreme factors in the body, activation of the sympathoadrenal system of the body occurs first of all [Karabaev A.G. 2020]. At the same time, the deterministic excitation of the higher sympathetic centers and neuroendocrine systems, an increase in the concentration of catecholamines, leads to the realization of two interrelated phenomena: activation of the adenylate cyclase





system, that is, energy conversion processes and sequential activation of the main processes of renewal of the lipid bi-layer of membranes, i.e. lipid peroxidation, the latter provides dystrophic changes at the cellular level. At the same time, the reproductive and regenerative processes in the cell remain in an unfavorable position. It is known that luteinizing hormone (LH), follicle-stimulating hormone (FSH) and testosterone are responsible for this kind of process [Hernandez-Hernandez J.M. et al., 2021; Bhasin, S. et al., 2018; Mulhall, J.P. et al., 2018; Mirone, V. et al., 2017].

Therefore, without knowing the sequence of dynamic changes in the components of the endocrine system responsible for the cellular reproductive process during the period of post-resuscitation disease, we cannot disclose the causes of changes in the cellular level and the disruption of reproductive standing at the cellular level.

An analysis of the literature shows that studies of endocrine system dysfunction during postrenimation disease, only some glands have been studied. That is, the hypothalamic-pituitary neurosecretory system, adrenal glands, thyroid gland, pancreas in the process of clinical death and post-resuscitation disease have been partially studied (Volkov A.V. et al., 2010; Zarechnova N.N., Slynko T.N., 2018; Yakimov I.A., Loginova E.S., 2017).

Certain measures are being taken in the world to create a healthcare system that provides a radical increase in the quality and effectiveness of medical care to the population, including early diagnosis, effective treatment, prevention and reduction of complications of various somatic diseases. Based on the identified tasks, it is advisable to conduct scientific research aimed at assessing disorders of the reproductive system in the post-resuscitation period.

The purpose of the study. It will reveal the reactivity of the reproductive system of males during clinical death.

Research objective

Assess the functional state of the velta and delta basophilic cells of the adenohipophysis, the content of follicle-stimulating hormone (FSH) lutinating hormone (LH), testosterone in the blood of intact rats.

Assess the functional state of the velta and delta basophilic cells of the adenohipophysis, the content of follicle-stimulating hormone (FSH) lutinating hormone (LH), testosterone in the blood of male rats when simulating clinical death duration of 10 minutes.

Object of research: In connection with the task, a study was conducted on 20 mongrel male rats weighing 160-170 g. Of these , it was: 10 intact, 10 experimental rats, in which a study of the reactivity of the reproductive system was conducted in modeling 10 clinical death.





Research methods. To achieve the goal and solve the problem, experimental, morphological, morphometric, histochemical, cytophotometric, enzyme immunoassay methods were used.

The results obtained and its discussion. In the study, β - and d-basophilic cells are scattered throughout the adenohypophysis. They are mainly defined around the vessels. β -basophilic adenocytes have a smaller size compared to d-basophilic adenocytes, in both basophilic adenocytes the glycoprotein is diffusely located, depending on functional activity. The cytoplasm volume in β -basophilic adenocytes is $592.1 \pm 17.9 \text{ mm}^3$, the cytoplasm volume of d-basophilic adenocytes is greater than that of β -basophilic adenocytes, which are within $812.0 \pm 31.3 \text{ mm}^3$. When determining the functional activity of β - and d – basophilic adenocytes, the cells are at the stage of different functional activity. At the same time, in β - and d-cells, the number of highly active cells is $10.6 \pm 0.4\%$ and $10.8 \pm 0.4\%$. The glycoprotein in them is located mainly in the pericaryon region. The preparation is mainly dominated by cells of moderate functional activity, the glycoprotein in them is scattered throughout the cytoplasm, diffusely and loosely, their number is equal to $70.6 \pm 0.4\%$ and $70.4 \pm 0.6\%$, and the number of cells of low functional activity, that is, with a densely located glycoprotein in the cytoplasm is determined within $18.8 \pm 0.4\%$ and $18.8 \pm 0.4\%$. The glycoprotein content is determined in the range of 155.8 ± 3.3 units and 156.0 ± 3.1 units (Fig. 1). The nuclei in β - and d-basophilic adenocytes are chromatic, chromatin in them is diffusely located throughout the nucleus, the volume of the nuclei is 127.2 ± 3.5 microns³ and 126.9 ± 3.5 microns³, the nucleoli in them are mainly located in the center of the nucleus, their volume is on average 0.88 ± 0.02 microns³ and 0.87 ± 0.02 microns³. The value of the nuclear-cytoplasmic ratio in β -basophilic adenocytes is equal to 0.215 ± 0.0009 , and in d-basophilic adenocytes, the value of the nuclear–cytoplasmic ratio is less, compared with β -basophilic adenocytes and is equal to 0.156 ± 0.001 .




Fig.1. Basophilic cells of the adenohypophysis in intact rats with a densely located glycoprotein. PAF coloring with azan coloring according to Haidengine.



Magnification: about 100x, about 15x. The numbers are: 1. d-basophilic adenocytes, 2. β -basophilic adenocytes.

The capillaries in the adenohypophysis have different sizes, they are slightly hyperemic. Their diameter is within 5.9 ± 0.05 microns.

The FSH content in the blood was 0.13 ± 0.015 mIU/ml. LH 0.43 ± 0.01 mIU/ml. testosterone content 5.04 ± 0.03 nmol/l

After the onset of clinical death, after 10 minutes, β - and d-basophil cells are clearly detected in the adenohypophysis, an increase in the number of depleted β - and d-basophil cells, that is, cells of high functional activity increased to $20.4 \pm 0.5\%$ ($P < 0.001$) and $21.2 \pm 0.5\%$ ($P < 0.001$) (Fig.3.2.6). The number of basophilic cells of moderate and low functional activity in β -cells decreased to $63.4 \pm 0.6\%$ and $16.2 \pm 1.0\%$ ($P < 0.01$), and in d-cells to $62.4 \pm 0.9\%$ ($P < 0.01$) and $16.4 \pm 0.8\%$ ($P < 0.05$) significantly compared with intact animals. In actively functioning basophilic cells, the glycoprotein is located mainly around the nucleus. In moderate functional activity, it is loose and scattered throughout the cytoplasm, and in basophilic cells of low functional activity, the glycoprotein is located intensively and densely throughout the cytoplasm. The cytoplasm volume in β -basophilic cells increased to 592.8 ± 19.6 mm³ ($P > 0.05$), and in d-basophilic cells to 831.6 ± 34.0 mm³ ($P > 0.05$) not significantly compared to intact animals.

The nucleus and nucleolus in the basophilic cells of the adenohypophysis are slightly swollen. Their volume in β -cells was increased to 127.7 ± 3.7 mm³ ($P > 0.05$) and 0.90 ± 0.02 mm³ ($P > 0.05$), and in d-basophilic cells to 128.4 ± 3.8 mm³ ($P > 0.05$) and 0.9 ± 0.02 mm³ ($P > 0.05$). The amount of glycoprotein in them decreased to 151.7 ± 1.6 units ($P > 0.05$) and 150.6 ± 1.6 units ($P < 0.05$).

The index of the nuclear-cytoplasmic ratio in β -basophil cells of the adenohypophysis is 0.215 ± 0.0009 ($P > 0.05$), and in d-basophil cells 0.154 ± 0.0002 ($P > 0.05$).




Fig.2. 10 minutes after the onset of clinical death, there is a further decrease in the amount of glycoprotein more in the d-basophil cells of the adenohypophysis. PAF



coloring with azan coloring according to Haidengine. Magnification: about 100x, about 15x. The numbers are: 1. β -basophilic adenocytes, 2. d- basophilic adenocytes.

Therefore, it can be said that the functional activity of d-basophil cells of the adenohipophysis is more pronounced compared to β -basophil cells of the adenohipophysis

At the same time, an increase in the FSH content to 0.17 ± 0.02 mIU/ml ($P > 0.05$), LH to 0.47 ± 0.01 mIU/ml ($P < 0.05$), the amount of testosterone to 6.15 ± 0.33 nmol/l ($P < 0.05$) was detected in the blood.

If we interpret the data with the data (Karabaeva A.G.2021), the reproductive system in intact rats is in the stage of quiet functioning. And an increase in the functional activity of the activity of d-basophil cells of the adenohipophysis, the content of the Lethinating hormone and the content of testosterone can indicate the activation of a compensatory mechanism aimed at supporting the reproductive activity of the body's cells to support the general and specific reactivity of the body against a subtle kind of stress factor.

Based on the obtained data, it is possible to draw the following conclusions.

- 1.The reproductive system in intatx rats is in a state of moderate functioning to reflect the synthesis and secretion of testosterone.
- 2.During the period of clinical death, there is an increase in the functional activity of d-basophil cells of the adenohipophysis, the content of the Lethinating hormone provided an increase in the secretion of testosterone to ensure the reproductive activity of the cells of the body.

Literature

1. Volkov A.V., Moroz V.V., Yezhova K.N., Zarzhetsky Yu.V. The role of sex steroids in the recovery period after clinical death (experimental study). General resuscitation.- 2010. 4(1):-p.1-18
2. Karabaev A. G. The relationship between the reactivity of the autonomic nervous system and the morphofunctional activity of basophilic cells of the adenohipophysis in the post-resuscitation period //Science and Peace. – 2020. – №. 3-1. – Pp. 55-61.
3. Zarechnova N.N., Slynko T.N. The effect of mountain hypoxia on the organs of the endocrine system with insufficiency of adrenal and pancreatic hormones // Bulletin of New Medical Technologies. Electronic edition-2018. No. 4.-pp.3-10.





4. Yakimov I. A., Loginova E. S. Analysis of changes in the level of thyroid hormones in certain types of death // Journal: Almanac of modern Science and Education. - 2017.No. 6.- pp.91-92
5. Bhasin, S. Testosterone therapy in men with hypogonadism / S. Bhasin, J.P. Brito, G.R. Cunningham et al. // An Endocrine Society clinical practice guideline. J. Clin. Endocrinol. Metab.– 2018. - Vol. 103 – P.1715–1744.
6. Hernández-Hernández, J.M. Kisspeptin Stimulatestion of Luteinizing Hormone (LH) during Postpartum Anestrus Continuous and Restricted Suckling / J.M. Hernández-Hernández Becerril-érez et al. // Animals (Basel). – 2021. – Vol.11 – P.1-8.
7. Mulhall, J.P. Evaluation and management of testosterone deficiency / J.P. Mulhall, L.W. Trost, R.E. Brannigan // AUA guideline. J. Urol. – 2018. - Vol. 200 – P.423–432.
8. Karabaev A.G. et al. Reactivity of the supraoptic, arcuate nucleus of the hypothalamus and the B-and D-basophilic cells of the adenohypophysis in the early postreanimation period //European Journal of Molecular & Clinical Medicine. – 2021. – Vol. 8. – No. 3. – pp. 954-957.
9. Karabaev A.G. Relationship between the reactivity of the autonomic nervous system and the morphofunctional activity of basophilic cells of the adenohypophysis in the post-resuscitation period. // Science and World International scientific journal- 2020. 3 (79). P.55-62.
10. Karabayev A. G., R. I. Isroilov. Morphofunctional Changes in Basophilic Cells of the denohypophysis during Post-resuscitation Disease // Journal of Advances in Medicine and Medical Research- 2020. 32 (8).p.130-135.
11. Mirone, V. European Association of Urology Position Statement on the role of the urologist in the management of male hypogonadism and testosterone therapy / V. Mirone, F. Debruyne, G. Dohle et al. // Eur. Urol. – 2017. - Vol. 72 – P.164–167.

